

Dear Parent / Guardian

Unfortunately a case of Head Lice has been reported in your child's class.

We are notifying you so you can check your own child's hair if you wish to. Attached is information provided by the National Assembly for Wales for guidance on how to detect and treat head lice.

Thank you.



These notes are intended for families and may be photocopied for circulation

What are head lice?

- Head lice are small insects (no larger than a sesame seed when fully grown) with moving legs. They live very close to the scalp.
- Nits are not the same as lice. They are the empty dead egg cases which stick to the hair, and look like specks of dandruff.
- You only have head lice if you find a moving louse (not a nit). Your head does not have to be itchy to have head lice.

How do you get head lice?

- Because of children's play activity and close contact, head lice are usually found on children, but can also spread to adults.
- Head lice infections are caught from close family and friends in the home and community, as well as at school.
- Head lice live in all types of hair - dirty or clean, short or long.
- Head lice walk from one head to another, when heads are close together - they do not fly, jump or swim.



How can you prevent head lice?

- Regular brushing and combing of the hair will help to spot lice early.
- The best way to stop the spread of head lice is for families to check their heads by regular wet combing using a detector comb.
- Human head lice are not found on animals or household pets and are not transmitted from pets to humans.

What are the side effects of head lice?

The reaction of individuals to louse bites can vary enormously from person to person. When you have head lice for the first time, you will probably not even realise they are there. However, after a short time (up to a few weeks), you start to itch, and react to the bites. Once you have had this initial reaction, you will experience itching and redness within 24-48 hours, each time you become re-infested. However, some individuals never exhibit symptoms. Itching from large numbers of bites may make the infested individual feel tired, irritable and feverish, hence the term 'feeling lousy'.

How do you detect head lice?

YOU WILL NEED: A DETECTOR COMB

(These can be bought from your pharmacist – ask for help if there are none on display.)

1. Wash the hair well and towel dry until damp, but not dripping.
2. Ensure that there is **good lighting** - daylight is best.
3. You could use a hand lens or magnifying glass (10x). Magnification may help detect nits and lice.
4. Comb the hair with an ordinary comb.
5. Using the detector comb, begin at the top of the head and making sure that the comb is touching the scalp, slowly draw the comb towards the ends of the hair. The use of conditioner may ease the process. Pay particular attention to the nape of the neck and behind the ears.
6. Check the teeth of the comb carefully for lice after each stroke. Wipe the comb on a white cloth/towel/kitchen roll to help you to see the lice.
7. Repeat steps (5) and (6) working your way around the head from the top of the scalp to the ends of the hair. If there are head lice, you will find one or more on the teeth of the comb. Once you have finished clean the comb under the tap with water.
8. If one member of the family has lice, check the rest of the family, including adults, as there is a good chance others are infected.





How do you treat head lice?

DO NOT TREAT UNLESS YOU ARE SURE YOU HAVE FOUND A MOVING LOUSE.

1. Check and treat all infected members of the family.
2. Ask your GP or Pharmacist which head lice treatment to use. Do not use head lice shampoo. The Pharmacist can advise on any evidence of resistance to treatment and therefore the most appropriate treatment to purchase. Also, the Pharmacist can advise on treatments suitable for people with asthma and other respiratory conditions.
3. Always read and follow the instructions in the treatment pack.
4. Apply the treatment carefully.

The usual methods are: Lotions and Liquids

In a well ventilated room (open a window), and away from all sources of heat (fires, cigarettes, hairdryers, lighters or matches).

- i. Apply the lotion or liquid to dry hair
- ii. Part the hair near the top of the head, put a few drops of the lotion or liquid on to the scalp and rub in. Part the hair again a bit further down the scalp and rub in some of the lotion or liquid. Do this again and again until the whole scalp is wet.

If you are treating long hair, ensure that you cover the hair down to where the ponytail band would be, UNLESS you are using Dimeticone (Hedrin) which should fully cover the hair. Ensure that you use enough lotion - do not dilute head lice treatments. Take care not to get the lotion or liquid in the eyes or on the face.

iii. Let the lotion or liquid dry on the hair naturally. Keep well away from naked flames, cigarettes or other lighted objects.

Do NOT use a hair dryer.

iv. Leave on the hair for the amount of time stated in the instructions. Then wash and rinse in the normal manner.

5. REPEAT THE ENTIRE TREATMENT SEVEN DAYS LATER, using a second bottle of the same lotion or liquid. This should kill any lice which may have hatched since the initial treatment.

6. Check the head two days after the second treatment. If you still find moving lice ask your pharmacist or GP for advice.

You should use at least one small bottle of lotion or liquid per head, more if the hair is thick.